arXiv:math/0501072v1 [math.CA] 5 Jan 2005

Asymptotic analysis of the Askey-scheme I: from Krawtchouk to Charlier

Diego Dominici [*]
Department of Mathematics
State University of New York at New Paltz
75 S. Manheim Blvd. Suite 9
New Paltz, NY 12561-2443
USA
Phone: (845) 257-2607
Fax: (845) 257-3571

February 8, 2008

---

[*] e-mail: dominicd@newpaltz.edu

Abstract

We analyze the Charlier polynomials $C_n(x)$ and their zeros asymptotically as $n \to \infty$. We obtain asymptotic approximations, using the limit relation between the Krawtchouk and Charlier polynomials, involving some special functions. We give numerical examples showing the accuracy of our formulas.

Keywords: Charlier polynomials, Askey-scheme, asymptotic analysis, orthogonal polynomials, hypergeometric polynomials, special functions.

MSC-class: 33C45 (Primary) 34E05, 33C10 (Secondary)

1 Introduction

The Charlier polynomials $C_n(x)$ [7] are defined by

$$C_n(x) = {}_2F_0\left(\begin{matrix} -n, -x \\ - \end{matrix} \middle| -\frac{1}{a} \right) \tag{1}$$

where $x \geq 0,\ n = 0, 1, \ldots$ and $a > 0$. They satisfy the discrete orthogonality condition [37]

$$\sum_{j=0}^{\infty} \frac{a^j}{j!} C_n(j) C_m(j) = a^{-n} e^a n! \delta_{nm}.$$

They are part of the Askey-scheme [18] of hypergeometric orthogonal polynomials:



where the arrows indicate limit relations between the polynomials.

The Charlier polynomials have applications in quantum mechanics [28], [31], [36], [39], difference equations [5], [23], teletraffic theory [16], [27], generating functions [4], [22], [26], and probability theory [3], [29], [30], [32]. The q-analogue of the Charlier polynomials were studied in [2], [8], [19] and [41]. The generalized Charlier polynomials were analyzed in [14], [17], [33], [34] and [40].

Asymptotics for the L^p-norms and information entropies of Charlier polynomials were derived in [21]. Bounds for their zeros were obtained in [20]. Asymptotic representations were established in [11] in terms of Hermite polynomials and in [24] in terms of Gamma functions. Some asymptotic estimates were computed in [15] from a representation of $C_n(x)$ in terms of Bell polynomials. An asymptotic formula when $x < 0$ was derived in [25] using probabilistic methods.

In [13], Goh studied the asymptotic behavior of $C_n(x)$ for large n using an approximation of the Plancharel-Rotach type. A uniform asymptotic expansion was derived in [6] using the saddle-point method. Asymptotic expansions were obtained in [10] from a second order linear differential satisfied by $C_n(x)$ in which a is the independent variable and x is a parameter.

In this paper we shall take a different approach and investigate the asymptotic behavior of $C_n(x)$ as $n \to \infty$, by using the limit relation between the Krawtchouk polynomials $K_n(x)$ defined by

$$K_n(x) = K_n(x, p, N) = {}_2F_1 \left(\begin{matrix} -n, -x \\ -N \end{matrix} \middle| \frac{1}{p} \right), \quad n = 0, 1, \ldots, N,\ 0 \le x \le N,\ 0 \le p \le 1 \tag{2}$$

and the Charlier polynomials, namely

$$\lim_{N \to \infty} K_n \left(x, \frac{a}{N}, N \right) = C_n(x). \tag{3}$$

We shall use the asymptotic expansions derived in [9] for the scaled Krawtchouk polynomials $k_n(x)$, with

$$k_n(x) = k_n(x, p, N) = (-p)^n \binom{N}{n} K_n(x, p, N). \tag{4}$$

A similar idea has been used in [12] and [38] to obtain asymptotic approximations of several orthogonal polynomials of the Askey-scheme in terms of Hermite and Laguerre polynomials.

2 Preliminaries

The following is the main result derived in [9].

Theorem 1 *As $N \to \infty$, $k_n(x, p, N)$ admits the following asymptotic approximations (see Figure1).*

1. $n = O(1)$, $0 \le y \le 1$, $y \not\approx p$.

$$k_n(x) \sim k_n^{(1)}(y) = \frac{\varepsilon^{-n}}{n!} (y - p)^n \tag{5}$$

where

$$\varepsilon = N^{-1}, \quad x = \frac{y}{\varepsilon}, \quad n = \frac{z}{\varepsilon} \quad 0 \le y, z \le 1.$$

2. $n = O(1)$, $y \approx p$, $y = p + \eta\sqrt{2pq\varepsilon}$, $\eta = O(1)$.

$$k_n(x) \sim k_n^{(2)}(\eta) = \frac{\varepsilon^{-\frac{n}{2}}}{n!} \left(\frac{pq}{2} \right)^{\frac{n}{2}} \mathrm{H}_n(\eta), \tag{6}$$

where $q = 1 - p$ and $\mathrm{H}_n(\eta)$ is the Hermite polynomial.



Figure 1: A sketch of the different asymptotic regions for $k_n(x)$.

3. $0 \le y < Y^{-}(z)$, $0 < z < p$, *where*

$$Y^{\pm}(z) = p + (q-p)\,z \pm 2zU_0, \quad U_0(z) = \sqrt{\frac{pq(1-z)}{z}}. \tag{7}$$

$$k_n(x) \sim k^{(3)}(y,z) = \frac{\sqrt{\varepsilon}}{\sqrt{2\pi}} \exp\left[\psi(y,z,U^{-})\varepsilon^{-1}\right] L(z,U^{-}), \tag{8}$$

with

$$\psi(y,z) = (z-1)\ln(U) + (1-y)\ln(U-p) + y\ln(U+q), \tag{9}$$

$$L(y,z) = \sqrt{\frac{(U-p)(U+q)}{z\left[U^2 - (U_0)^2\right]}} \tag{10}$$

and

$$U^{\pm}(y,z) = -\frac{1}{2}\left(\frac{p-y}{z} + q - p\right) \pm \frac{1}{2}\sqrt{\left(\frac{p-y}{z} + q - p\right)^2 - 4\,(U_0)^2}. \tag{11}$$

4. $Y^{+}(z) < y \le 1$, $0 < z < q$.

$$k_n(x) \sim k^{(4)}(y,z) = \frac{\sqrt{\varepsilon}}{\sqrt{2\pi}} \exp\left[\psi(y,z,U^{+})\varepsilon^{-1}\right] L(z,U^{+}). \tag{12}$$

5. $x = O(1)$, $p < z < 1$.

$$\begin{aligned} k_n(x) \sim k^{(5)}(x,z) = {} & \frac{\sqrt{\varepsilon}}{\sqrt{2\pi}\sqrt{z\,(1-z)}} \cos(\pi x) \left(\frac{z-p}{p}\right)^x \exp\left[\phi_0(z)\varepsilon^{-1}\right] \\ & -\frac{\varepsilon}{\pi}\frac{x}{z-p}\Gamma(x)\sin(\pi x)\left(\frac{q\varepsilon}{z-p}\right)^x \exp\left[\frac{(z-1)\ln(q) + \pi \mathrm{i} z}{\varepsilon}\right] \end{aligned} \tag{13}$$

where

$$\phi_0(z) = (z-1)\ln(1-z) - z\ln(z) + z\ln(-p)$$

and $\Gamma(x)$ *is the Gamma function.*

6. $x = O(1)$, $z \approx p$, $z = p - u\sqrt{pq\varepsilon}$, $u = O(1)$.

$$\begin{aligned} k_n(x) \sim k^{(6)}(x,u) = {} & \frac{\sqrt{\varepsilon}}{\sqrt{2\pi pq}}\left[\sqrt{\frac{q\varepsilon}{p}}\right]^x \mathrm{D}_x(u) \\ & \times \exp\left[\frac{\pi \mathrm{i} p - q\ln(q)}{\varepsilon} + \frac{u\sqrt{pq}\pi\mathrm{i} - u\sqrt{pq}\ln(q)}{\sqrt{\varepsilon}} - \frac{u^2}{4}\right], \end{aligned} \tag{14}$$

where $\mathrm{D}_x(u)$ *is the parabolic cylinder function.*

7. $0 \ll y < Y^{-}(z)$, $p < z < 1$.

$$k_n(x) \sim k^{(7)}(y,z) = \exp\left(\frac{\pi \mathrm{i} y}{\varepsilon}\right)\left[\cos\left(\frac{\pi y}{\varepsilon}\right) k^{(4)}(y,z) + 2\mathrm{i}\sin\left(\frac{\pi y}{\varepsilon}\right) k^{(3)}(y,z)\right] \tag{15}$$

8. $y \approx Y^{-}(z)$, $0 < z < p$, $y = Y^{-}(z) - \beta\varepsilon^{2/3}$, $\beta = O(1)$.

$$k_n(x) \sim k^{(8)}(\beta,z) = \varepsilon^{\frac{1}{3}} \exp\left[\psi_0(z)\varepsilon^{-1} + \ln\left(\frac{U_0+p}{U_0-q}\right)\beta\varepsilon^{-\frac{1}{3}}\right] \mathrm{Ai}\left(\Theta^{\frac{2}{3}}\beta\right)\frac{\Theta^{-\frac{1}{3}}}{\sqrt{zU_0}}, \tag{16}$$

where

$$\psi_0(z) = z\pi\mathrm{i} + (z-1)\ln(U_0) + Y^{-}(z)\ln(U_0-q) + \left[1 - Y^{-}(z)\right]\ln(U_0+p), \tag{17}$$

$$\Theta(z) = \sqrt{\frac{U_0}{z}}\frac{1}{(U_0+p)(U_0-q)} \tag{18}$$

and $\mathrm{Ai}(\cdot)$ *is the Airy function.*

9. $y \approx Y^{-}(z)$, $p < z < 1$, $\quad y = Y^{-}(z) - \beta\varepsilon^{\frac{2}{3}}$, $\beta = O(1)$.

$$k_n(x) \sim k^{(9)}(\beta,z) = \varepsilon^{\frac{1}{3}} \exp\left[\psi_0(z)\varepsilon^{-1} + \ln\left(\frac{U_0+p}{U_0-q}\right)\beta\varepsilon^{-\frac{1}{3}}\right] \tag{19}$$

$$\times \frac{1}{2}\frac{\vartheta^{-\frac{1}{3}}}{\sqrt{zU_0}}\left[\lambda^{+}(\beta,z)\mathrm{Ai}\left(\vartheta^{\frac{2}{3}}\beta\right) + \mathrm{i}\lambda^{-}(\beta,z)\mathrm{Bi}\left(\vartheta^{\frac{2}{3}}\beta\right)\right],$$

where $\vartheta(z) = -\Theta(z)$,

$$\lambda^{\pm}(\beta,z) = \exp\left\{2\pi\mathrm{i}\left[Y^{-}(z) - \beta\varepsilon^{\frac{2}{3}}\right]\varepsilon^{-1}\right\} \pm 1. \tag{20}$$

and $\mathrm{Ai}(\cdot), \mathrm{Bi}(\cdot)$ *are the Airy functions.*

10. $Y^{-}(z) < y < Y^{+}(z)$, $0 < z < 1$.

$$k_n(x) \sim k^{(10)}(y,z) = k^{(3)}(y,z) + k^{(4)}(y,z). \tag{21}$$

11. $y \approx Y^{+}(z)$, $0 < z < q$, $\quad y = Y^{+}(z) + \alpha\varepsilon^{\frac{2}{3}}$, $\quad \alpha = O(1)$.

$$k_n(x) \sim k^{(11)}(\alpha,z) = \varepsilon^{\frac{1}{3}} \exp\left[\psi_1(z)\varepsilon^{-1} + \ln\left(\frac{U_0+q}{U_0-p}\right)\alpha\varepsilon^{-\frac{1}{3}}\right] \mathrm{Ai}\left[(\Theta_1)^{\frac{2}{3}}\alpha\right]\frac{(\Theta_1)^{-\frac{1}{3}}}{\sqrt{zU_0}}, \tag{22}$$

where

$$\psi_1(z) = (z-1)\ln(U_0) + Y^{+}(z)\ln(U_0+q) + \left[1 - Y^{+}(z)\right]\ln(U_0-p), \tag{23}$$

$$\Theta_1(z) = \sqrt{\frac{U_0}{z}}\frac{1}{(U_0-p)(U_0+q)} \tag{24}$$

In order to obtain the corresponding asymptotic expansions for $K_n(x)$, we need to derive asymptotic formulas for $(-p)^n \binom{N}{n}$ in the different regions of Theorem 1.

The following lemma follows immediately from Stirling's formula [1]

$$\Gamma(x) \sim \sqrt{\frac{2\pi}{x}} x^x e^{-x}, \quad x \to \infty. \tag{25}$$

Lemma 2 *As $N \to \infty$, we have the following asymptotic approximations:*

1.
$$(-p)^n \binom{N}{n} \sim \frac{1}{n!} (-p)^n \varepsilon^{-n}, \quad n = O(1). \tag{26}$$

2.
$$(-p)^n \binom{N}{n} \sim \frac{\sqrt{\varepsilon}}{\sqrt{2\pi}\sqrt{z(1-z)}} \exp\left[\phi(z)\varepsilon^{-1}\right], \quad n = z\varepsilon^{-1} \tag{27}$$

   where
$$\phi(z) = z\ln(p) + z\pi\mathrm{i} + (z-1)\ln(1-z) - z\ln(z). \tag{28}$$

3.
$$(-p)^n \binom{N}{n} \sim \frac{\sqrt{\varepsilon}}{\sqrt{2\pi pq}} \exp\left[\phi_1(u)\right], \quad n = p\varepsilon^{-1} - u\sqrt{\frac{pq}{\varepsilon}}, \quad u = O(1) \tag{29}$$

   where
$$\phi_1(u) = \left[\pi\mathrm{i}p - q\ln(q)\right]\varepsilon^{-1} - u\sqrt{pq}\left[\ln(q) + \pi\mathrm{i}\right]\varepsilon^{-\frac{1}{2}} - \frac{1}{2}u^2. \tag{30}$$

3 Limit analysis

Setting
$$y = x\varepsilon, \quad z = n\varepsilon, \quad q = 1-p, \quad p = a\varepsilon \tag{31}$$
in (7) and letting $\varepsilon \to 0$, we obtain

$$Y^{\pm}(z) \to X^{\pm}(a,n) = \left(\sqrt{n} \pm \sqrt{a}\right)^2, \quad U_0(z) \to \sqrt{\frac{a}{n}}. \tag{32}$$

Hence, the eleven regions of Theorem 1 transform into the following regions (see Figure 2).

1. Region I

   From (5) and (26) we have for $n = O(1)$
$$K_n(x) \sim \left(1 - \frac{y}{p}\right)^n. \tag{33}$$



Figure 2: A sketch of the different asymptotic regions for $C_n(x)$.

Thus,

$$C_n(x) \simeq \left(1 - \frac{x}{a}\right)^n . \tag{34}$$

The formula above is exact for $n = 0, 1$ and in the limit as $a \to \infty$ we have

$$\lim_{a\to\infty} C_n(av) = (1 - v)^n = {}_1F_0\left(\begin{array}{c} -n \\ - \end{array} \middle| \, v \right).$$

2. Region II

   From (6) and (26) we have for $x = p\varepsilon^{-1} + \eta\sqrt{\frac{2pq}{\varepsilon}}, \quad \eta = O(1)$

   $$K_n(x) \sim (-1)^n \left(\frac{\varepsilon q}{2p}\right)^{\frac{n}{2}} \mathrm{H}_n\left(\eta\right). \tag{35}$$

   Hence,

   $$C_n(x) \simeq (-1)^n \left(2a\right)^{-\frac{n}{2}} \mathrm{H}_n\left(\eta\right), \quad x = a + \eta\sqrt{2a}. \tag{36}$$

   Equation (36) is exact for $n = 0, 1$ and in the limit as $a \to \infty$ we have

   $$\lim_{a\to\infty} \left(-1\right)^n \left(2a\right)^{\frac{n}{2}} C_n(a + \eta\sqrt{2a}) = \mathrm{H}_n\left(\eta\right)$$

   which is equation 2.12.1 in.

3. Region III

   From (8), (27) and (28) we have for $0 \leq y < Y^-(z), \quad 0 < z < p$

   $$K_n(x) \sim K^{(3)}(y, z) = \exp\left[\frac{\psi\left(y, z, U^-\right) - \phi(z)}{\varepsilon}\right] G\left(z, U^-\right), \tag{37}$$

   where

   $$G(z, U) = \sqrt{\frac{(1 - z)(U - p)(U + q)}{U^2 - Uo^2}}. \tag{38}$$

   From (9), (11) and (38) we obtain

   $$\frac{\psi\left(y, z, U^-\right) - \phi(z)}{\varepsilon} \to \Psi_3(x),$$

   where

   $$\Psi_3(x) = x \ln\left(\frac{a + x - n + \Delta}{2a}\right) + n \ln\left(\frac{a - x + n + \Delta}{2a}\right) + \frac{1}{2}\left(a - x - n + \Delta\right) \tag{39}$$

   and

   $$G\left(z, U^-\right) \to L_3(x) \equiv \sqrt{\frac{a - x - n + \Delta}{2\Delta}}, \tag{40}$$



Figure 3: A comparison of $C_n(x)$ (solid curve) and $F_3(x)$ (ooo) for $n = 30$ with $a = 50.165184$.

for $0 \leq x < X^-, \quad 0 < n < a$, with

$$\Delta(a, n, x) = \sqrt{a^2 - 2a(x+n) + (x-n)^2}. \tag{41}$$

Thus,

$$C_n(x) \sim F_3(x) = \exp\left[\Psi_3(x)\right] L_3(x), \quad 0 \leq x < X^-, \quad 0 < n < a. \tag{42}$$

Figure 3 shows the accuracy of the approximation (42) with $n = 30$ and $a = 50.165184$ in the range $-3 < x < X^-$.

Remark 3 *Although we have not shown proof of it, the approximation (42) is in fact also valid for* $x < 0$ *and* $n \geq 0$ *(see Figure 4).*

4. Region IV



Figure 4: A comparison of $C_n(x)$ (solid curve) and $F_3(x)$ (ooo) for $n = 30$ with $a = 2.165184$.

From (12), (27) and (28) we have for $Y^+(z) < y \leq 1, \quad 0 < z < q$

$$K_n(x) \sim K^{(4)}(y,z) = \exp\left[\frac{\psi(y,z,U^+) - \phi(z)}{\varepsilon}\right] G\left(z,U^+\right). \tag{43}$$

From (9), (11) and (38) we obtain

$$\frac{\psi(y,z,U^+) - \phi(z)}{\varepsilon} \to \Psi_4(x) - n\pi \mathrm{i},$$

where

$$\Psi_4(x) = x\ln\left(\frac{a+x-n-\Delta}{2a}\right) + n\ln\left(\frac{x-a-n+\Delta}{2a}\right) + \frac{1}{2}(a-x-n+\Delta) \tag{44}$$

and

$$G\left(z,U^-\right) \to L_4(x) \equiv \sqrt{\frac{x-a+n+\Delta}{2\Delta}}, \tag{45}$$

for $X^+ < x$. Therefore,

$$C_n(x) \sim F_4(x) = (-1)^n \exp\left[\Psi_4(x)\right] L_4(x), \quad X^+ < x. \tag{46}$$

Figure 5 shows the accuracy of the approximation (46) with $n = 30$ and $a = 2.165184$ in the range $X^+ < x < \infty$.

5. Region V

From (13), (27) and (28) we have for $x = O(1), \quad p < z < 1$

$$\begin{aligned} K_n(x) \sim \exp\left[x\ln\left(\frac{z}{p}-1\right)\right]\cos(\pi x) - \frac{\sqrt{\varepsilon}}{z-p}\sqrt{\frac{2}{\pi}z(1-z)}\Gamma(x+1)\sin(\pi x) \\ \times \exp\left[\frac{(1-z)\ln\left(\frac{1-z}{q}\right) + z\ln\left(\frac{z}{p}\right)}{\varepsilon} + x\ln\left(\frac{\varepsilon q}{z-p}\right)\right]. \end{aligned} \tag{47}$$

Hence,

$$\begin{aligned} C_n(x) \sim F_5(x) = \exp\left[x\ln\left(\frac{n}{a}-1\right)\right]\cos(\pi x) - \sqrt{n}\sqrt{\frac{2}{\pi}}\Gamma(x+1)\sin(\pi x) \\ \times \exp\left[n\ln\left(\frac{n}{a}\right) - (x+1)\ln(n-a) + a - n\right], \quad x \approx 0, \quad n > a. \end{aligned} \tag{48}$$

Figure 6 shows the accuracy of the approximation (48) with $n = 30$, $a = 2.165184$ and $x \approx 0$.



Figure 5: A comparison of $C_n(x)$ (solid curve) and $F_4(x)$ (ooo) for $n = 30$ with $a = 2.165184$.



Figure 6: A comparison of $C_n(x)$ (solid curve) and $F_5(x)$ (ooo) for $n = 30$ with $a = 2.165184$.



Figure 7: A comparison of $C_n(x)$ (solid curve) and $F_6(x)$ (ooo) for $n = 30$ with $a = 30.165184$.

6. Region VI

   From (14) and (29)-(30) we have for $x = O(1)$, $\quad z = p - u\sqrt{pq\varepsilon}$, $\quad u = O(1)$

   $$K_n(x) \sim \exp\left[\frac{x}{2}\ln\left(\frac{q\varepsilon}{p}\right) + \frac{u^2}{4}\right] \mathrm{D}_x(u). \tag{49}$$

   Thus,

   $$C_n(x) \sim F_6(x) = \exp\left[-\frac{x}{2}\ln(a) + \frac{u^2}{4}\right] \mathrm{D}_x(u), \quad x \approx 0, \quad n = a - u\sqrt{a}. \tag{50}$$

   Figure 7 shows the accuracy of the approximation (50) with $n = 30$, $a = 30.165184$ ($n \approx a$) and $x \approx 0$.

7. Region VII

From (15), (37) and (43) we have for $0 \ll y < Y^{-}(z)$, $p < z < 1$

$$K_n(x) \sim \exp\left(\frac{\pi \mathrm{i} y}{\varepsilon}\right)\left[\cos\left(\frac{\pi y}{\varepsilon}\right) K^{(4)}(y,z) + 2\mathrm{i}\sin\left(\frac{\pi y}{\varepsilon}\right) K^{(3)}(y,z)\right]. \tag{51}$$

Therefore

$$C_n(x) \sim \exp(\pi \mathrm{i} x)\left[\cos(\pi x)\, C_n^{(4)}(x) + 2\mathrm{i}\sin(\pi x)\, C_n^{(3)}(x)\right]$$

for $0 \ll x < X^{-},\ n > a$, which we can rewrite as

$$\begin{aligned} C_n(x) \sim F_7(x) = {} & \exp\left[x\ln\left(\frac{n-a-x+\Delta}{2a}\right) + n\ln\left(\frac{a+n-x-\Delta}{2a}\right) + \frac{1}{2}(a-x-n+\Delta)\right] \\ & \times \cos(\pi x)\sqrt{\frac{x+n-a+\Delta}{2\Delta}} - 2\sin(\pi x)\sqrt{\frac{x+n-a-\Delta}{2\Delta}} \\ & \times \exp\left[x\ln\left(\frac{n-a-x-\Delta}{2a}\right) + n\ln\left(\frac{a+n-x+\Delta}{2a}\right) + \frac{1}{2}(a-x-n+\Delta)\right]. \end{aligned} \tag{52}$$

Figure 8 shows the accuracy of the approximation (52) with $n = 30$ and $a = 2.165184$ in the range $0 \ll x < X^{-}$.

8. Region VIII

From (16), (27) and (28) we have for $y \approx Y^{-}(z)$, $0 < z < p$, $y = Y^{-}(z) - \beta\varepsilon^{2/3}$, $\beta = O(1)$

$$\begin{aligned} K_n(x) \sim {} & \varepsilon^{-\frac{1}{6}} \exp\left[\frac{\psi_0(z) - \phi(z)}{\varepsilon} + \ln\left(\frac{U_0 + p}{U_0 - q}\right)\beta\varepsilon^{-\frac{1}{3}}\right] \\ & \times \sqrt{2\pi}\left[\frac{z(1-z)}{pq}\right]^{\frac{1}{4}} \mathrm{Ai}\left(\Theta^{\frac{2}{3}}\beta\right)\Theta^{-\frac{1}{3}}. \end{aligned} \tag{53}$$

Since

$$\beta = \left[Y^{-}(z) - y\right]\varepsilon^{-\frac{2}{3}} \tag{54}$$

we have from (32)

$$\beta\varepsilon^{-\frac{1}{3}} \to X^{-} - x. \tag{55}$$

From (17), (28) and (55) we get

$$\frac{\psi_0(z) - \phi(z)}{\varepsilon} + \ln\left(\frac{U_0 + p}{U_0 - q}\right)\beta\varepsilon^{-\frac{1}{3}} \to \frac{1}{2} n \ln\left(\frac{n}{a}\right) + x\ln\left(1 - \sqrt{\frac{n}{a}}\right) + \sqrt{an} - \sqrt{n}.$$

From (18) and (55) we obtain

$$\varepsilon^{-\frac{1}{6}}\sqrt{2\pi}\left[\frac{z(1-z)}{pq}\right]^{\frac{1}{4}}\Theta^{-\frac{1}{3}} \to \sqrt{2\pi}\left(\frac{n}{a}\right)^{\frac{1}{6}}\left(\sqrt{a} - \sqrt{n}\right)^{\frac{1}{3}} \tag{56}$$



Figure 8: A sketch of $C_n(x)/F_7(x)$ for $n = 30$ with $a = 2.165184$. The vertical lines are due to the discontinuities at $x \simeq 13, 14$ and 15.

and

$$\Theta^{\frac{2}{3}}\beta \to \left(\frac{n}{a}\right)^{\frac{1}{6}} \frac{(X^{-}-x)}{\left(\sqrt{a}-\sqrt{n}\right)^{\frac{2}{3}}}. \tag{57}$$

Therefore,

$$\begin{aligned} C_n(x) \sim F_8(x) &= \sqrt{2\pi}\left(\frac{n}{a}\right)^{\frac{1}{6}}\left(\sqrt{a}-\sqrt{n}\right)^{\frac{1}{3}} \operatorname{Ai}\left[\left(\frac{n}{a}\right)^{\frac{1}{6}} \frac{(X^{-}-x)}{\left(\sqrt{a}-\sqrt{n}\right)^{\frac{2}{3}}}\right] \\ &\times \exp\left[\frac{1}{2}n\ln\left(\frac{n}{a}\right)+x\ln\left(1-\sqrt{\frac{n}{a}}\right)+\sqrt{an}-\sqrt{n}\right] \end{aligned} \tag{58}$$

for $x \approx X^{-},\ 0<n<a.$

9. Region IX

From (19), (20), (27) and (28) we have for $y \approx Y^{-}(z),\ p<z<1,\quad y=Y^{-}(z)-\beta\varepsilon^{\frac{2}{3}},\ \beta=O(1)$

$$\begin{aligned} K_n(x) &\sim \varepsilon^{-\frac{1}{6}}\exp\left[\frac{\psi_0(z)-\phi(z)}{\varepsilon}+\ln\left(\frac{U_0+p}{U_0-q}\right)\beta\varepsilon^{-\frac{1}{3}}\right] \\ &\times \sqrt{2\pi}\left[\frac{z\,(1-z)}{pq}\right]^{\frac{1}{4}}\vartheta^{-\frac{1}{3}}\left[\lambda^{+}(\beta,z)\operatorname{Ai}\left(\vartheta^{\frac{2}{3}}\beta\right)+\mathrm{i}\lambda^{-}(\beta,z)\operatorname{Bi}\left(\vartheta^{\frac{2}{3}}\beta\right)\right], \end{aligned}$$

which can be written as

$$\begin{aligned} K_n(x) &\sim \varepsilon^{-\frac{1}{6}}\sqrt{2\pi}\exp\left[\frac{(z-1)\ln\left(\frac{U_0}{1-z}\right)+Y^{-}\ln\left(q-U_0\right)+\left(1-Y^{-}\right)\ln\left(p+U_0\right)+z\ln\left(\frac{z}{p}\right)}{\varepsilon}\right] \\ &\times \exp\left[\ln\left(\frac{U_0+p}{q-U_0}\right)\beta\varepsilon^{-\frac{1}{3}}\right]\left[\frac{z\,(1-z)}{pq}\right]^{\frac{1}{4}}\vartheta^{-\frac{1}{3}}\left[\cos\left(\pi x\right)\operatorname{Ai}\left(\vartheta^{\frac{2}{3}}\beta\right)-\sin(\pi x)\operatorname{Bi}\left(\vartheta^{\frac{2}{3}}\beta\right)\right]. \end{aligned} \tag{59}$$

Using (56) and (57) in (59) with $\vartheta=-\Theta$, we have

$$\begin{aligned} C_n(x) \sim F_9(x) &= \sqrt{2\pi}\left(\frac{n}{a}\right)^{\frac{1}{6}}\left(\sqrt{n}-\sqrt{a}\right)^{\frac{1}{3}}\exp\left[\frac{1}{2}n\ln\left(\frac{n}{a}\right)+x\ln\left(\sqrt{\frac{n}{a}}-1\right)+\sqrt{an}-n\right] \\ &\times\left\{\cos\left(\pi x\right)\operatorname{Ai}\left[\left(\frac{n}{a}\right)^{\frac{1}{6}}\frac{(X^{-}-x)}{\left(\sqrt{n}-\sqrt{a}\right)^{\frac{2}{3}}}\right]-\sin(\pi x)\operatorname{Bi}\left[\left(\frac{n}{a}\right)^{\frac{1}{6}}\frac{(X^{-}-x)}{\left(\sqrt{n}-\sqrt{a}\right)^{\frac{2}{3}}}\right]\right\} \end{aligned} \tag{60}$$

for $x \approx X^{-},\ n>a.$



Figure 9: A comparison of $C_n(x)$ (solid curve) and $F_{10}(x)$ (ooo) for $n = 30$ with $a = 2.165184$.

10. Region X

    From (21), (37) and (43) we have for $Y^-(z) < y < Y^+(z),\ 0 < z < 1$

$$K_n(x) \sim K^{(3)}(y, z) + K^{(4)}(y, z). \tag{61}$$

Thus,

$$C_n(x) \sim F_{10}(x) = F_3(x) + F_4(x), \quad X^- < x < X^+. \tag{62}$$

Figure 9 shows the accuracy of the approximation (62) with $n = 30$, $a = 2.165184$ in the range $X^- < x < X^+$.

11. Region XI

    From (22), (27) and (28) we have for $y \approx Y^+(z)$, $0 < z < q$, $\quad y = Y^+(z) + \alpha\varepsilon^{\frac{2}{3}}, \quad \alpha =$

$O(1)$,

$$K_n(x) \sim \varepsilon^{-\frac{1}{6}} \exp\left[\frac{\psi_1(z)-\phi(z)}{\varepsilon} + \ln\left(\frac{U_0+q}{U_0-p}\right) \alpha\varepsilon^{-\frac{1}{3}}\right] \tag{63}$$
$$\times \sqrt{2\pi}\left[\frac{z\,(1-z)}{pq}\right]^{\frac{1}{4}} \mathrm{Ai}\left[(\Theta_1)^{\frac{2}{3}}\beta\right](\Theta_1)^{-\frac{1}{3}}.$$

Since

$$\alpha = \left[y - Y^+(z)\right]\varepsilon^{-\frac{2}{3}} \tag{64}$$

we have from (32)

$$\alpha\varepsilon^{-\frac{1}{3}} \to x - X^+. \tag{65}$$

From (23), (28) and (65) we get

$$\frac{\psi_1(z)-\phi(z)}{\varepsilon} + \ln\left(\frac{U_0+q}{U_0-p}\right)\alpha\varepsilon^{-\frac{1}{3}} \to \frac{1}{2}n\ln\left(\frac{n}{a}\right) + x\ln\left(1+\sqrt{\frac{n}{a}}\right) - \sqrt{an} - \sqrt{n} - n\pi\mathrm{i}.$$

From (24) and (65) we obtain

$$\varepsilon^{-\frac{1}{6}}\sqrt{2\pi}\left[\frac{z\,(1-z)}{pq}\right]^{\frac{1}{4}}(\Theta_1)^{-\frac{1}{3}} \to \sqrt{2\pi}\left(\frac{n}{a}\right)^{\frac{1}{6}}\left(\sqrt{a}+\sqrt{n}\right)^{\frac{1}{3}} \tag{66}$$

and

$$(\Theta_1)^{\frac{2}{3}}\alpha \to \left(\frac{n}{a}\right)^{\frac{1}{6}} \frac{(x-X^+)}{(\sqrt{a}+\sqrt{n})^{\frac{2}{3}}}. \tag{67}$$

Therefore,

$$C_n(x) \sim F_{11}(x) = \sqrt{2\pi}\left(\frac{n}{a}\right)^{\frac{1}{6}}\left(\sqrt{a}+\sqrt{n}\right)^{\frac{1}{3}} \mathrm{Ai}\left[\left(\frac{n}{a}\right)^{\frac{1}{6}}\frac{(x-X^+)}{(\sqrt{a}+\sqrt{n})^{\frac{2}{3}}}\right] \tag{68}$$
$$\times(-1)^n \exp\left[\frac{1}{2}n\ln\left(\frac{n}{a}\right) + x\ln\left(1+\sqrt{\frac{n}{a}}\right) - \sqrt{an} - \sqrt{n}\right]$$

for $x \approx X^+$.

4 Comparison with previous results

We shall now compare our results with those obtained previously in [6] and [13].

1. Region VII: $0 \le x < X^-$, $n > a$.

   Setting $x = un$, with

$$u = O(1), \quad 0 \le u < 1 - 2\sqrt{\frac{a}{n}} + \frac{a}{n} < 1$$

in (52) we have, as $n \to \infty$

$$F_7(x) \sim g_7(u) = \frac{\cos(un\pi)}{\sqrt{1-u}} \exp\left\{ \left[u \ln\left(\frac{n}{a}\right) + (u-1)\ln(1-u) - u \right] n + \frac{au}{u-1} \right\} \tag{69}$$
$$-2\sin(un\pi)\sqrt{\frac{u}{1-u}} \exp\left\{ \left[\ln\left(\frac{n}{a}\right) + (1-u)\ln(1-u) + u\ln(u) - 1 \right] n + \frac{a}{1-u} \right\}.$$

The second term of equation (69) is the same as the equation before (5.3) in [6] and equation (84) in [13]. However, the first term is absent in previous works, although it is necessary in the asymptotic approximation, especially when $u \simeq 0, 1, 2, \ldots$.

2. Region IX: $x \approx X^-, \ n > a$.

   We now set $x = X^- + tn^{\frac{1}{6}}, \ t = O(1)$ in (60) and obtain, as $n \to \infty$

$$F_9(x) \sim g_9(t) = \sqrt{2\pi} a^{-\frac{1}{6}} n^{\frac{1}{3}} \exp\left[\frac{1}{2}\left(X^- + tn^{\frac{1}{6}} + n \right) \ln\left(\frac{n}{a}\right) - n + \frac{3}{2}a \right] \tag{70}$$
$$\times \left\{ \cos\left[\left(X^- + tn^{\frac{1}{6}} \right) \pi \right] \mathrm{Ai}\left(-ta^{-\frac{1}{6}} \right) - \sin\left[\left(X^- + tn^{\frac{1}{6}} \right) \pi \right] \mathrm{Bi}\left(-ta^{-\frac{1}{6}} \right) \right\}.$$

   Equation (70) agrees with equation (5.13) in [6] and equation (51) in [13].

3. Region X: $X^- < x < X^+$.

   Setting $x = n + a + 2\sin(\theta)\sqrt{an}$, with $-\frac{\pi}{2} < \theta < \frac{\pi}{2}$ in (42) we have, as $n \to \infty$

$$F_3(x) \sim g_3(\theta) = (-1)^n \frac{a^{-\frac{1}{4}} n^{\frac{1}{4}}}{\sqrt{2\cos(\theta)}} \exp\left\{ \left[\ln\left(\frac{n}{a}\right) - 1 \right] n + \frac{\pi}{4}\mathrm{i} \right\}$$
$$\times \exp\left\{ \sqrt{an} \left[\sin(\theta)\ln\left(\frac{n}{a}\right) - \sin(\theta)(2\theta - \pi)\mathrm{i} - 2\cos(\theta)\mathrm{i} \right] \right\} \tag{71}$$
$$\times \exp\left\{ a\left[1 - \frac{1}{2}\cos(2\theta) + \frac{1}{2}\ln\left(\frac{n}{a}\right) - \frac{1}{2}\sin(2\theta)\mathrm{i} - \theta\mathrm{i} + \frac{\pi}{2}\mathrm{i} \right] \right\}.$$

   Similarly from (46) we get

$$F_4(x) \sim g_4(\theta) = (-1)^n \frac{a^{-\frac{1}{4}} n^{\frac{1}{4}}}{\sqrt{2\cos(\theta)}} \exp\left\{ \left[\ln\left(\frac{n}{a}\right) - 1 \right] n - \frac{\pi}{4}\mathrm{i} \right\}$$
$$\times \exp\left\{ \sqrt{an} \left[\sin(\theta)\ln\left(\frac{n}{a}\right) + \sin(\theta)(2\theta - \pi)\mathrm{i} + 2\cos(\theta)\mathrm{i} \right] \right\} \tag{72}$$
$$\times \exp\left\{ a\left[1 - \frac{1}{2}\cos(2\theta) + \frac{1}{2}\ln\left(\frac{n}{a}\right) + \frac{1}{2}\sin(2\theta)\mathrm{i} + \theta\mathrm{i} - \frac{\pi}{2}\mathrm{i} \right] \right\}.$$

Using (71) and (72) in (62) we have

$$\begin{aligned} F_{10}(x) \sim g_{10}(\theta) &= (-1)^n \frac{\sqrt{2} a^{-\frac{1}{4}} n^{\frac{1}{4}}}{\sqrt{\cos(\theta)}} \exp\left\{ \left[\ln\left(\frac{n}{a}\right) - 1 \right] n \right\} \\ &\times \exp\left\{ \sqrt{an} \left[\sin(\theta) \ln\left(\frac{n}{a}\right) \right] + a \left[1 - \frac{1}{2}\cos(2\theta) + \frac{1}{2}\ln\left(\frac{n}{a}\right) \right] \right\} \\ &\times \cos\left\{ \sqrt{an} \left[\sin(\theta)(2\theta - \pi) + 2\cos(\theta) \right] + a \left[\frac{1}{2}\sin(2\theta) + \theta - \frac{\pi}{2} \right] - \frac{\pi}{4} \right\}. \end{aligned} \tag{73}$$

Equation (73) is equivalent to equation (44) in [13].

4. Region XI: $x \approx X^+$.

   We now set $x = X^+ + sn^{\frac{1}{6}},\ s = O(1)$ in (68) and obtain, as $n \to \infty$

$$F_{11}(x) \sim g_{11}(s) = \sqrt{2\pi} a^{-\frac{1}{6}} n^{\frac{1}{3}} \exp\left[\frac{1}{2}\left(X^+ + sn^{\frac{1}{6}} + n \right) \ln\left(\frac{n}{a}\right) - n + \frac{3}{2}a \right] \mathrm{Ai}\left(sa^{-\frac{1}{6}} \right). \tag{74}$$

   Equation (74) is equation (5.12) in [6] and equation (30) in [13].

5 Zeros

Using the formulas from the previous sections we can obtain approximations to the zeros of the Charlier polynomials.

1. $x \simeq 0,\ n > a$.

   The first zero is exponentially small. From (48) we have, as $x \to 0$

$$C_5(x) \sim 1 + \left[\ln\left(\frac{n}{a} - 1\right) - \frac{\sqrt{2\pi n}}{n - a} a^{-n} n^n e^{a-n} \right] x.$$

   Solving for x we obtain

$$x_0 \simeq \left[\ln\left(\frac{n}{a} - 1\right) - \frac{\sqrt{2\pi n}}{n - a} a^{-n} n^n e^{a-n} \right]^{-1} \sim \frac{e^{n-a} a^n n^{-n}}{\sqrt{2\pi n}}, \quad n \to \infty \tag{75}$$

   where x_0 denotes the smallest zero.

2. $0 < x < X^-,\ n > a$.

   In this range of x, the zeros are exponentially close to $1, 2, \ldots, \lfloor X^- \rfloor$. Using

$$t = \frac{x - n - a + 2\sqrt{an}}{n^{\frac{1}{6}}}$$

and the asymptotic formulas [35]

$$\operatorname{Ai}(x) \sim \frac{\exp\left[-\frac{2}{3}x^{\frac{3}{2}}\right]}{2\sqrt{\pi}x^{\frac{1}{4}}}, \quad x \to \infty$$

$$\operatorname{Bi}(x) \sim \frac{\exp\left[\frac{2}{3}x^{\frac{3}{2}}\right]}{\sqrt{\pi}x^{\frac{1}{4}}}, \quad x \to \infty$$

we have, as $n \to \infty$

$$\frac{\operatorname{Ai}\left(-ta^{\frac{1}{6}}\right)}{\operatorname{Bi}\left(-ta^{\frac{1}{6}}\right)} \sim \frac{1}{2}\exp\left[-\frac{4}{3}a^{-\frac{1}{4}}n^{-\frac{1}{4}}\left(X^{-}-x\right)^{\frac{3}{2}}\right]. \tag{76}$$

Using (76) in (70) we have

$$g_9(t) \simeq 0 \Leftrightarrow \frac{1}{2}\exp\left[-\frac{4}{3}a^{-\frac{1}{4}}n^{-\frac{1}{4}}\left(X^{-}-x\right)^{\frac{3}{2}}\right] \simeq \tan\left(\pi x\right).$$

Since $x_j \simeq j,\ j = 1, 2, \ldots, \lfloor X^{-} \rfloor$, we get

$$\frac{1}{2}\exp\left[-\frac{4}{3}a^{-\frac{1}{4}}n^{-\frac{1}{4}}\left(X^{-}-j\right)^{\frac{3}{2}}\right] \simeq \pi\left(x_j - j\right)$$

which we can solve to obtain

$$x_j \simeq j + \frac{\pi}{2}\exp\left[-\frac{4}{3}a^{-\frac{1}{4}}n^{-\frac{1}{4}}\left(X^{-}-j\right)^{\frac{3}{2}}\right], \quad j = 1, 2, \ldots, \lfloor X^{-} \rfloor. \tag{77}$$

3. $X^{-} < x < X^{+}$.

   Finally, the non-trivial zeros of the Charlier polynomials can be approximated using (73). We have $g_{10}(\theta) = 0$ if and only if

$$\cos\left\{\sqrt{an}\left[\sin(\theta)(2\theta-\pi) + 2\cos(\theta)\right] + a\left[\frac{1}{2}\sin(2\theta) + \theta - \frac{\pi}{2}\right] - \frac{\pi}{4}\right\} = 0$$

or equivalently if

$$\sqrt{an}\left[\sin(\theta)(2\theta-\pi) + 2\cos(\theta)\right] + a\left[\frac{1}{2}\sin(2\theta) + \theta - \frac{\pi}{2}\right] - \frac{\pi}{4} = \frac{\pi}{2} + \pi l,\ l \in \mathbb{Z}$$

or

$$\sqrt{an}\left[\sin(\theta)(2\theta-\pi) + 2\cos(\theta)\right] + a\left[\frac{1}{2}\sin(2\theta) + \theta - \frac{\pi}{2}\right] - \frac{3\pi}{4} - \pi l = 0, \tag{78}$$

with $-\frac{\pi}{2} < \theta < \frac{\pi}{2}$. Recalling that

$$x = n + a + 2\sin(\theta)\sqrt{an}, \tag{79}$$

we see that the condition $X^- < x < X^+$ implies

$$0 \leq l \leq 2\sqrt{an} - a - \frac{3}{4}. \tag{80}$$

Equation (78) cannot be solved exactly. However, it can be easily solved numerically to any desired accuracy and using (79) gives very good approximations for the nontrivial zeros.

In Table 1 we computed the exact and approximate zeros of $C_{25}(x)$ with $a = 2.16564899$ using (75), (77) and (78)-(80).

Conclusion 4 *We analyzed the asymptotic behavior of the Charlier polynomials in the range* $0 \leq x$ *as* $n \to \infty$. *We also obtained approximations for their zeros. We intend to extend our method to the other polynomials of the Askey-scheme to obtain asymptotic expansions of them.*

References

[1] M. Abramowitz and I. A. Stegun, editors. *Handbook of mathematical functions with formulas, graphs, and mathematical tables.* Dover Publications Inc., New York, 1992. Reprint of the 1972 edition.

[2] N. Asai. Integral transform and Segal-Bargmann representation associated to q-Charlier polynomials. In *Quantum information, IV (Nagoya, 2001)*, pages 39–48. World Sci. Publishing, River Edge, NJ, 2002.

[3] A. D. Barbour. Asymptotic expansions in the Poisson limit theorem. *Ann. Probab.*, 15(2):748–766, 1987.

[4] N. Barik. Some theorems on generating functions for Charlier polynomials. *J. Pure Math.*, 3:111–114, 1983.

[5] H. Bavinck and R. Koekoek. On a difference equation for generalizations of Charlier polynomials. *J. Approx. Theory*, 81(2):195–206, 1995.

[6] R. Bo and R. Wong. Uniform asymptotic expansion of Charlier polynomials. *Methods Appl. Anal.*, 1(3):294–313, 1994.

[7] C. Charlier. Über die Darstellung willkürlicher Funktionen. *Ark. Mat. Astron. Fys.*, 2(20):1–35, 1906.

Table 1: Comparison of the exact and approximate zeros of $C_{25}(x)$ with $a = 2.16564899$.

l	x (exact)	x (approximate)
–	$0.41229323 \times 10^{-16}$	$0.41549221 \times 10^{-16}$
–	1.0000000	1.0000000
–	2.0000000	2.0000000
–	3.0000000	3.0000001
–	4.0000000	4.0000009
–	5.0000001	5.0000073
–	6.0000015	6.0000507
–	7.0000227	7.0003063
–	8.0002574	8.0015785
–	9.0021153	9.0068260
–	10.012329	10.024179
–	11.050278	11.067497
–	12.147166	12.137242
11	13.330606	13.334295
10	14.615276	14.560867
9	16.007976	15.899727
8	17.514470	17.350792
7	19.142918	18.921714
6	20.905595	20.626110
5	22.820702	22.484600
4	24.915443	24.527911
3	27.232157	26.803591
2	29.842164	29.391394
1	32.883964	32.446240
0	36.717784	36.379078

[8] A. de Médicis, D. Stanton, and D. White. The combinatorics of q-Charlier polynomials. *J. Combin. Theory Ser. A*, 69(1):87–114, 1995.

[9] D. E. Dominici. Asymptotic analysis of the Krawtchouk polynomials by the WKB method. Preprint, arXiv: math.CA/0501042, 2004.

[10] T. M. Dunster. Uniform asymptotic expansions for Charlier polynomials. *J. Approx. Theory*, 112(1):93–133, 2001.

[11] C. Ferreira, J. L. López, and E. Mainar. Asymptotic approximations of orthogonal polynomials. In *Seventh Zaragoza-Pau Conference on Applied Mathematics and Statistics (Spanish) (Jaca, 2001)*, volume 27 of *Monogr. Semin. Mat. García Galdeano*, pages 275–280. Univ. Zaragoza, Zaragoza, 2003.

[12] C. Ferreira, J. L. Lopez, and E. Mainar. Asymptotic relations in the Askey scheme for hypergeometric orthogonal polynomials. *Adv. in Appl. Math.*, 31(1):61–85, 2003.

[13] W. M. Y. Goh. Plancherel-Rotach asymptotics for the Charlier polynomials. *Constr. Approx.*, 14(2):151–168, 1998.

[14] M. N. Hounkonnou, C. Hounga, and A. Ronveaux. Discrete semi-classical orthogonal polynomials: generalized Charlier. *J. Comput. Appl. Math.*, 114(2):361–366, 2000.

[15] L. C. Hsu. Certain asymptotic expansions for Laguerre polynomials and Charlier polynomials. *Approx. Theory Appl. (N.S.)*, 11(1):94–104, 1995.

[16] D. L. Jagerman. Nonstationary blocking in telephone traffic. *Bell System Tech. J.*, 54:625–661, 1975.

[17] G. C. Jain and R. P. Gupta. On a class of polynomials and associated probabilities. *Utilitas Math.*, 7:363–381, 1975.

[18] R. Koekoek and R. F. Swarttouw. The Askey-scheme of hypergeometric orthogonal polynomials and its q-analogue. Technical Report 98-17, Delft University of Technology, 1998. http://aw.twi.tudelft.nl∼koekoek/askey/.

[19] H. T. Koelink. Yet another basic analogue of Graf's addition formula. *J. Comput. Appl. Math.*, 68(1-2):209–220, 1996.

[20] I. Krasikov. Bounds for zeros of the Charlier polynomials. *Methods Appl. Anal.*, 9(4):599–610, 2002.

[21] L. Larsson-Cohn. L^p-norms and information entropies of Charlier polynomials. *J. Approx. Theory*, 117(1):152–178, 2002.

[22] P. A. Lee. Some generating functions involving the Charlier polynomials. *Nanta Math.*, 8(1):83–87, 1975.

[23] J. Letessier. Some results on co-recursive associated Meixner and Charlier polynomials. *J. Comput. Appl. Math.*, 103(2):323–335, 1999.

[24] J. L. López and N. M. Temme. Convergent asymptotic expansions of Charlier, Laguerre and Jacobi polynomials. *Proc. Roy. Soc. Edinburgh Sect. A*, 134(3):537–555, 2004.

[25] M. Maejima and W. Van Assche. Probabilistic proofs of asymptotic formulas for some classical polynomials. *Math. Proc. Cambridge Philos. Soc.*, 97(3):499–510, 1985.

[26] E. B. McBride. *Obtaining generating functions.* Springer Tracts in Natural Philosophy, Vol. 21. Springer-Verlag, New York, 1971.

[27] M. L. Mehta and E. A. van Doorn. Inequalities for Charlier polynomials with application to teletraffic theory. *J. Math. Anal. Appl.*, 133(2):449–460, 1988.

[28] J. Negro and L. M. Nieto. Symmetries of the wave equation in a uniform lattice. *J. Phys. A*, 29(5):1107–1114, 1996.

[29] N. Privault. Multiple stochastic integral expansions of arbitrary Poisson jump times functionals. *Statist. Probab. Lett.*, 43(2):179–188, 1999.

[30] B. Roos. Poisson approximation of multivariate Poisson mixtures. *J. Appl. Probab.*, 40(2):376–390, 2003.

[31] A. Ruffing, J. Lorenz, and K. Ziegler. Difference ladder operators for a harmonic Schrödinger oscillator using unitary linear lattices. In *Proceedings of the Sixth International Symposium on Orthogonal Polynomials, Special Functions and their Applications (Rome, 2001)*, volume 153, pages 395–410, 2003.

[32] W. Schoutens. Lévy-Sheffer and IID-Sheffer polynomials with applications to stochastic integrals. In *Proceedings of the VIIIth Symposium on Orthogonal Polynomials and Their Applications (Seville, 1997)*, volume 99, pages 365–372, 1998.

[33] B. Şefik. Coherent structures in nonlinear dynamical systems. Method of the random point functions. In *Nonlinear evolution equations and dynamical systems (Baia Verde, 1991)*, pages 385–394. World Sci. Publishing, River Edge, NJ, 1992.

[34] R. M. Shreshtha. On generalised Charlier polynomials. *Nepali Math. Sci. Rep.*, 7(2):65–69, 1982.

[35] J. Spanier and K. B. Oldham. *An Atlas of Functions.* Hemisphere Pub. Corp., 1987.

[36] F. H. Szafraniec. Charlier polynomials and translational invariance in the quantum harmonic oscillator. *Math. Nachr.*, 241:163–169, 2002.

[37] G. Szegő. *Orthogonal polynomials.* American Mathematical Society, Providence, R.I., fourth edition, 1975. American Mathematical Society, Colloquium Publications, Vol. XXIII.

[38] N. M. Temme and J. L. López. The Askey scheme for hypergeometric orthogonal polynomials viewed from asymptotic analysis. In *Proceedings of the Fifth International Symposium on Orthogonal Polynomials, Special Functions and their Applications (Patras, 1999)*, volume 133, pages 623–633, 2001.

[39] T. T. Truong. On a class of inhomogeneous Ising quantum chains. *J. Phys. A*, 28(24):7089–7096, 1995.

[40] W. Van Assche and M. Foupouagnigni. Analysis of non-linear recurrence relations for the recurrence coefficients of generalized Charlier polynomials. *J. Nonlinear Math. Phys.*, 10(suppl. 2):231–237, 2003.

[41] J. Zeng. The q-Stirling numbers, continued fractions and the q-Charlier and q-Laguerre polynomials. *J. Comput. Appl. Math.*, 57(3):413–424, 1995.